

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



B4

(11)

EP 0 435 081 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
12.08.1998 Bulletin 1998/33

(51) Int Cl.⁶: C07D 249/08, A61K 31/41

(21) Application number: 90123986.3

(22) Date of filing: 12.12.1990

(54) **Triazole derivatives**

Triazololderivate

Dérivés de triazole

(84) Designated Contracting States:
BE CH DE ES FR GB IT LI NL SE

(30) Priority: 25.12.1989 JP 335946/89
09.10.1990 JP 270783/90

(43) Date of publication of application:
03.07.1991 Bulletin 1991/27

(73) Proprietor: **SS PHARMACEUTICAL CO., LTD.**
Chuo-ku, Tokyo 103 (JP)

(72) Inventors:

- Tokizawa, Minoru
Narita-shi, Chiba (JP)
- Kanamaru, Yoshihiko
Narita-shi, Chiba (JP)
- Matsumoto, Masaru
Inba-gun, Chiba (JP)
- Asaka, Takemitsu
Narita-shi, Chiba (JP)
- Matsuda, Hideaki
Abiko-shi, Chiba (JP)

- Kuraishi, Tadayuki
Chiba-shi, Chiba (JP)
- Maebashi, Kazunori
Narashino-shi, Chiba (JP)
- Taido, Naokata
Funabashi-shi, Chiba (JP)
- Kawahara, Ryuichi
Ichikawa-shi, Chiba (JP)

(74) Representative:
Wächtershäuser, Günter, Prof. Dr.
Patentanwalt,
Tal 29
80331 München (DE)

(56) References cited:
EP-A- 61 835 EP-A- 95 828
EP-A- 332 387

Remarks:

The file contains technical information submitted
after the application was filed and not included in this
specification

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 0 435 081 B1

Description

BACKGROUND OF THE INVENTION5 Field of the Invention:

This invention relates to novel triazole derivatives and salts thereof possessing excellent antimycotic activity and to a composition for curing mycoses comprising the same.

10 Description of the Background Arts:

Mycosis is classified into skin diseases typified by *trichophytosis*, *psoriasis*, skin *candidasis*, and the like and deep-seated mycoses which include mycotic meningitis, mycotic infectious diseases of respiratory organs, fungemia, urinary tract mycosis, and the like.


15 Of these, patients of deep-seated mycoses are increasing, since they cannot be cured by conventional antibiotics or chemotherapeutic agents. Development of a drug efficacious for the treatment of these diseases has therefore been desired.

Very few drugs for curing deep mycoses are known. Only ketoconazole, fluconazole, and the like among azole compounds have been known as drugs for curing deep-seated mycoses. Their effects, however, are not necessarily satisfactory. Triazole compounds are known from EP-A-0 332 387; EP-A-0 095 828; EP-A-0 061 835.

SUMMARY OF THE INVENTION

25 In view of this situation, the present inventors have synthesized a number of triazole compounds and studied their antimycotic activities, and have found that triazole derivatives having the following formula (I) exhibited excellent antimycotic activity.

Accordingly, an object of the present invention is to provide a triazole derivative of formula (I),



45 wherein R¹ and R² represent a hydrogen atom or a C₁₋₆-alkyl group, or R¹ and R² together form an ethylene, trimethylene or tetramethylene group, R³ represents a C₁₋₆-alkyl group, and n denotes an integer of 0 to 2, provided that R¹ and R² are not both a hydrogen atom at the same time, and a salt thereof; and a composition for curing mycoses comprising such a triazole derivative or a salt thereof.

Other objects, features and advantages of the invention will hereinafter become more readily apparent from the following description.

50 DETAILED DESCRIPTION OF THE INVENTION AND PREFERRED EMBODIMENTS


A C₁₋₆ alkyl group represented by R¹, R², and R³ in formula (I) is a linear or branched alkyl group having 1 to 6 carbon atoms. Specific examples are methyl-, ethyl-, n-propyl-, isopropyl-, n-butyl-, isobutyl-, sec-butyl-, tert-butyl-, n-pentyl-, and n-hexyl groups.

Hydrochloride, nitrate, hydrobromide, sulfate, p-toluenesulfonate, methanesulfonate, fumarate, maleate, succinate, lactate, and the like are given as examples of the salts of the compound of formula (I).

There are optical isomers due to the asymmetric carbon atom and sulfur atom for the compound of formula (I). All

of racemates, optical active isomers, and diastereomers are included in this invention.

Triazole derivatives of formula (I) and their salts can be prepared, for example, according to the following reactions.



wherein R¹, R², and R³ have the same meanings as defined above.

First, compound (II) is epoxylated and reacted with triazole to produce sulfide compound (I-1). The sulfide compound (I-1) is then oxidized to obtain sulfinyl compound (I-2) or sulfonyl compound (I-3). Each step is described in more detail below.

The reaction to produce sulfide compound (I-1) from compound (II) is preferably carried out by using 1 to 2 moles of epoxylation agent, e.g. trimethylsulfoxonium iodide, and 1 to 3 moles of triazole for 1 mole of compound (II) in the presence of 2 to 5 moles of an alkali compound and by heating the mixture at room temperature to 100°C for 1 to 30 hours. Sodium hydroxide, potassium hydroxide, barium hydroxide, sodium carbonate, potassium carbonate, or the like can be used as an alkali. Of these, alkali hydroxides are particularly preferable. An alcohol such as methanol, ethanol, propanol, isopropanol, butanol, isobutanol, sec-butanol, tert-butanol, or the like is used as a solvent.

After completion of the reaction, the organic solvent is evaporated, ice-cooled water is added to the residue, and the residue is extracted with a solvent such as chloroform or the like. The target compound can be obtained by drying the chloroform layer, evaporating the solvent, and purifying by silica gel column chromatography or the like means.

When the sulfide compound (I-1) has different substituents; i.e., different R¹ and R², two types of diastereomers are produced; A isomer fractionated first by silica gel chromatography and B isomer which elutes later from the column.

The reaction for producing sulfinyl compound (I-2) from sulfide compound (I-1) is carried out by using 1 mole of oxidizing agent for 1 mole of sulfide compound (I-1) and by reacting at -10°C to room temperature for 0.1 to 2 hours. Organic peroxide, manganese dioxide, chromic acid, hydrogen peroxide, or the like can be used as the oxide for the reaction, with particularly preferable oxides being organic peroxides. A halogenated hydrocarbon is used as a preferable solvent. After completion of the reaction, the reaction mixture is washed with an alkali carbonate and then water, dried, distilled for removal of the solvent, and subjected to silica gel column chromatography for purification to obtain the target compound.


When the sulfinyl compound (I-2) has the same substituents R¹ and R², two types of diastereomers are produced; a isomer fractionated first by silica gel chromatography and b isomer which elutes later from the column. When R¹ and R² are different from each other, both a and b isomers are produced for each one of A and B isomers of sulfide compound (I-1).

The reaction for producing sulfonyl compound (I-3) from sulfide compound (I-1) is carried out by using 2 to 5 mol s

of oxidizing agent for 1 mole of sulfide compound (I-1) at room temperature to 50°C for 1 to 30 hours. The same oxidizing agent, solvent, and purification method as those for producing sulfinyl compound (I-2) can be employed.

Two types of diastereomers are present for sulfonyl compound (I-3), when the substituents R¹ and R² are different. When either one of A or B isomer of sulfide compound (I-1) is used, only one of the diastereomers of sulfonyl compound (I-3) is produced.

When the carbon atom which have a hydroxyl group substituent in the compound of formula (I) is the only asymmetric atom, there are two optical isomers present due to this asymmetric atom. Such optical isomers can be produced, for example, by the following reactions.



In the reaction scheme, R^1 , R^2 , and R^3 have the same meanings as defined above, and compounds (IV-1) and (IV-2) are diastereomers.

In the above reaction scheme, compound (II) is first reacted with optically active sulfinyl compound (III) to produce a mixture of diastereomers (IV-1) and (IV-2). Only one of the two diastereomers (IV-1) and (IV-2) is separated from the mixture and oxidized to produce sulfonyl compound (V). The sulfonyl compound (V) is then converted into compound (VI) by the reaction with acetyl chloride and sodium iodide. After epoxylation, the compound (VI) is reacted with triazole to obtain optically active compound (I-4).

Each of the above reactions are described in more detail.

Diastereomers (IV-1) and (IV-2) are produced by reacting 1 to 1.2 moles of optically active sulfinyl compound (III) for 1 mole of compound (II) in the presence of 1 to 1.4 moles of a base at 0°C to room temperature for 0.5 to 5 hours. Ether, tetrahydrofuran, dioxane, n-hexane, or the like is used as a solvent. As a base, lithium diisopropylamide, butyl lithium, potassium t-butoxide, sodium amide, sodium hydride, calcium hydride, or the like can be used. Of these, a particularly preferable base is lithium diisopropylamide. In order to collect the target compound from the mixture containing two diastereomers, the reaction mixture is treated with ammonium chloride or the like, extracted with an organic solvent, and purified by silica gel column chromatography or the like. Purified diastereomer (VI-1) is thus obtained.

Diastereomer (VI-1) is then dissolved into a solvent such as chloroform, dichloromethane, or the like and to the solution 2 moles of an oxidizing agent is added for 1 mole of diastereomer (VI-1). The mixture is stirred at -10°C to room temperature for 0.5 to 5 hours to obtain sulfonyl compound (V). Hydrogen peroxide, perbenzoic acid, m-chloroperoxybenzoic acid, or the like is used as an oxidizing agent. After the reaction, the reaction mixture is washed with an alkali and purified by silica gel column or the like to obtain sulfonyl compound (V).


Sulfonyl compound (V) thus obtained is dissolved into a solvent such as acetone and to the solution 1 to 3 moles of acetyl chloride and 1 to 3 moles of sodium iodide for 1 mole of sulfonyl compound (V) is added. The mixture is stirred at -10°C to room temperature for 0.5 to 5 hours to obtain compound (VI). After the reaction, the reaction mixture is extracted with an organic solvent and purified by silica gel column or the like to obtain purified compound (VI).

Compound (VI) is dissolved into chloroform, dichloromethane, or the like, and to the solution is added 2 to 10 moles of triethyloxonium tetrafluoroborate for 1 mole of compound (VI). The mixture is stirred at 0°C to room temperature for 10 to 30 hours to produce epoxylated compound (VII). After an addition of an alkali, the crude product is stirred at 0°C to room temperature for 0.5 to 5 hours. The organic layer is submitted to silica gel column or the like to obtain purified epoxylated compound (VII).

Epoxylated compound (VII) is dissolved into a solvent such as N,N-dimethylformamide, dimethylsulfoxide, acetonitrile, t-butanol, or the like, and to the solution are added 1 to 10 moles of triazole and 1 to 10 moles of an alkali for 1 mole of compound (VII). The mixture is stirred at 0°C to refluxing temperature for 1 to 30 hours to obtain the target optical active compound (I-4). Sodium hydroxide, potassium hydroxide, sodium carbonate, potassium carbonate, sodium bicarbonate, potassium bicarbonate, or the like can be used as an alkali. Purified optical active compound (I-4) can be obtained by extracting the reaction mixture with an organic solvent, evaporating the solvent, and recrystallizing from a suitable solvent.

The raw material compound (II) can easily be prepared according to the following reaction scheme (a) or (b).

Reaction Scheme (a):




5

10

15

20



25

wherein X represents a halogen atom, R^1 , R^2 , and R^3 represent a lower alkyl group.

According to the reaction scheme (a), 4-(trifluoromethyl)phenacyl halide (VIII) is converted into the sulfide derivative (IX) by the treatment with a lower alkyl mercaptan in the presence of an alkali, followed by the reaction with a lower alkyl halide to produce monoalkyl derivative (II-1). If required, monoalkyl derivative (II-1) is further reacted with the lower alkyl halide to produce dialkyl derivative (II-2).

30

Reaction Scheme (b):


35


40

45

50

55





wherein X^1 and X^2 represent a halogen atom, R^1 , R^2 , and R^3 have the same meanings as defined previously.

According to the reaction scheme (b), 4-(trifluoromethyl)benzaldehyde (X) is converted into alcohol compound (XI) by the Grignard reaction, oxidized into ketone compound (XII), and successively halogenized into compound (XIII). Compound (XIII) is then reacted with mercaptan to obtain compound (II).

Antimycotic activity of compounds of formula (I) of the present invention is described by way of the following experiment.

Groups of ICR male mice (age: 4 weeks), each group consisting of 5 mice, were subjected to the experiment. 3.3×10^6 *Candida albicans* were injected into caudal vein of each mouse. Starting from 1 hour after the injection, test compounds dissolved in 0.5 ml of 0.06 N hydrochloric acid solution to which 0.2% TO-10M (Trade mark: product of Nikko Chemicals Co.) was added were orally administered once each day, four times in total. The mice were observed for 14 days after the injection of *Candida albicans* to compare the average survival days of the mice to which the test compounds were administered with the average survival days of the control group. The indicator T/C (%) were determined according to the following equation.

$$T/C (\%) = \frac{\text{Average survival days of the mice to which the test compounds were administered}}{\text{Average survival days of the control}}$$

The results are presented in Table 1, in which Compound Nos. are the same as those given in Table 2 hereinafter.

TABLE 1

Compound No.	Dose (mg/kg)	Survived animals/tested animals	T/C (%)
7	5	5/5	412
	2.5	3/5	394
	1.25	0/5	235
10	5	3/5	406
	2.5	2/5	394
	1.25	1/5	306
	0.625	1/5	235
16	5	5/5	412
	2.5	3/5	400
	1.25	2/5	371
	0.625	2/5	341

TABLE 1 (continued)

Compound No.	Dose (mg/kg)	Survived animals/tested animals	T/C (%)
Fluconazole (Comparative compound)	5	2/5	300
	2.5	1/5	312
	1.25	1/5	271
	0.625	0/5	141
Control	-	0/5	100

As demonstrated by the above results, in the oral administration compounds of formula (I) the present invention exhibited a superior survival rate; the *in vivo* antimycotic effect, over fluconazole, the comparative compound.

When the compound of formula (I) of this invention or a salt thereof is used as an antimycotic agent, it is preferably orally administered at a dose of 50 to 500 mg per day or parenterally at a dose of 10 to 200 mg per day, depending on the weight, the age, the sex, the physical conditions, or the symptom of the patient or the way of administration.

The compound of formula (I) can be formed into various preparations, such as tablets, granules, powders, capsules, suspensions, injections, suppositories, or other forms for external application, according to known methods. When solid preparations are produced, the compound of formula (I) is mixed with excipients, and as required, with binders, disintegrators, lubricants, coloring agents, sweetening agents, flavoring agents, fillers, coating agents, sugar-coating agents, and the like, and formed into preparations such as tablets, granules, powders, capsules, suppositories, or the like according to known methods. The compound of the present invention can be also prepared into preparations for injection by dissolving, suspending, or emulsifying it into an aqueous medium such as distilled water, or by making it into powder which is dissolvable when it is injected. Subcutaneous, intravenous, intra-arterial, intra-portal, intraperitoneal, or intramuscular injection is applicable.

As discussed above, the compound of the present invention possesses high antimycotic activity. In addition, it is easily absorbed by living bodies, exhibiting its superior bioavailability. Thus, the compound is useful for curing mycoses or preventing infectious diseases in human and animals.

Other features of the invention will become apparent in the course of the following description of the exemplary embodiments which are given for illustration of the invention and are not intended to be limiting thereof.

EXAMPLES

Reference Example 1

To 13.2 g of 2-bromo-4'-(trifluoromethyl)acetophenone dissolved in 40 ml of methanol was added under ice-cooling 34.6 g of a 15% aqueous solution of methyl mercaptan sodium salt. After the reaction at room temperature for 1 hour with stirring, the resultant reaction mixture was poured into ice-cooled water and extracted with dichloromethane. The organic layer was washed with water and dried over anhydrous magnesium sulfate, followed by evaporation of the solvent to obtain 11.3 g of oily product of 2-methylthio-4'-(trifluoromethyl)acetophenone (yield: 97.7%).

NMR(CDCl₃) δ : 2.13 (s, 3H), 3.77 (s, 2H), 7.74(d, 2H), 8.17(d, 2H)

Reference Example 2

To 5.00 g of 2-methylthio-4'-(trifluoromethyl)acetophenone dissolved in 50 ml of tetrahydrofuran was added under ice-cooling 0.855 g of a 60% aqueous solution of sodium hydride, and the mixture was reacted at room temperature with stirring until no bubbles were produced. After an addition of 3.34 g of methyl iodide, the mixture was stirred at room temperature for a further 2 hours. The resultant reaction mixture was poured into ice-cooled water and extracted with dichloromethane. The organic layer was washed with water and dried over anhydrous magnesium sulfate. After evaporation of the solvent, the residue was purified by silica gel column chromatography using n-hexane to obtain 2.16 g of an oily product of 2-methylthio-4'-(trifluoromethyl)propiofenone (yield: 40.8%).

NMR(CDCl₃) δ : 1.55(d, 3H), 1.94(s, 3H), 4.30(q, 1H), 7.72(d, 2H), 8.11(d, 2H)

Reference Example 3

To 2.02 g of 2-methylthio-4'-(trifluoromethyl)-propiofenone dissolved in 20 ml of tetrahydrofuran was added under ice-cooling 0.33 g of a 60% aqueous solution of sodium hydride, and the mixture was reacted at room temperature

with stirring until no bubbles were produced. After an addition of 1.27 g of methyl iodide, the mixture was stirred at room temperature for a further 2 hours. The resultant reaction mixture was poured into ice-cooled water and extracted with dichloromethane. The organic layer was washed with water and dried over anhydrous magnesium sulfate. After evaporation of the solvent, the residue was purified by silica gel column chromatography using n-hexane to obtain 1.27 g of an oily product of 2-methylthio-4'-(trifluoromethyl)isobutyrophenone (yield: 59.5%).

NMR(CDCl₃)δ: 1.54(s, 6H), 2.01(s, 3H), 7.66(d, 2H), 8.11(d, 2H)

Reference Example 4

To a tetrahydrofuran solution of Grignard reagent prepared from 5.35 g of magnesium and 24.6 g of isopropyl bromide was added under ice-cooling 17.4 g of 4-(trifluoromethyl)benzaldehyde dissolved in tetrahydrofuran, and the mixture was stirred for 30 minutes. The resultant reaction mixture was treated with a saturated NH₄Cl aqueous solution and extracted with ether. The organic layer was washed with water and dried over anhydrous magnesium sulfate. After evaporation of the solvent, the residue was purified by silica gel column chromatography using n-hexane/ dichloromethane to obtain 8.02 g of an oily product of 1-(trifluoromethyl)phenyl-2-methylpropane-1-ol (yield: 36.8%).

NMR(CDCl₃)δ: 0.84, 0.96(d, 3H for each), 1.7-2.1 (m, 1H), 4.47(d, 1H), 7.43, 7.58(d, 2H for each)

Reference Example 5

To 8 g of 1-(trifluoromethyl)phenyl-2-methylpropane-1-ol dissolved in dichloromethane was added 15.8 g of pyridinium chlorochromate, and the mixture was stirred at room temperature for 3 hours. After separating the deposited undissolvable substance by filtration, the filtrate was concentrated and the residue was purified by silica gel column chromatography to obtain 7.9 g of an oily product of 4-(trifluoromethyl)isobutyrophenone (yield: 100%).

NMR(CDCl₃)δ: 1.24(d, 6H), 3.4-3.7(m, 1H), 7.71, 8.04 (d, 2H for each)

Reference Example 6

To 1 g of 4-(trifluoromethyl)isobutyrophenone dissolved in acetic acid was added 0.81 g of bromine at 40°C, and the mixture was stirred for 3 hours. The reaction mixture was poured into ice-cooled water and extracted with ether. The organic layer was washed with a 5% sodium bicarbonate aqueous solution and then with water, dried over anhydrous magnesium sulfate, and the solvent was evaporated to obtain 1.36 g of an oily product of 2-bromo-4'-(trifluoromethyl)isobutyrophenone (yield: 100%).

NMR(CDCl₃)δ: 2.03(s, 6H), 7.71, 8.19(d, 2H for each)

Reference Example 7

To 1 g of 2-bromo-4'-(trifluoromethyl)isobutyrophenone dissolved in methanol was added under ice-cooling 1.74 g of 15% solution of methyl mercaptan sodium salt, and the mixture was stirred for 30 minutes. The reaction mixture was poured into ice-cooled water and extracted with dichloromethane. The organic layer was washed with water, dried over anhydrous magnesium sulfate, and the solvent was evaporated. The residue was purified by silica gel column chromatography using n-hexane/dichloromethane to obtain 0.61 g of an oily product of 2-methylthio-4'-(trifluoromethyl)isobutyrophenone (yield: 68.7%).

NMR(CDCl₃)δ: 1.54(s, 6H), 2.01(s, 3H), 7.66, 8.11 (d, 2H)

Example 1

Into 50 ml of t-butanol were dissolved 4.97 g of 2-methylthio-4'-(trifluoromethyl)propiophenone, 8.05 g of trimethylsulfoxonium iodide, 5.61 g of potassium hydroxide, and 3.18 g of 1,2,4-triazole, and the mixture was stirred at 80°C for 7 hours. After the reaction, the solvent was evaporated, water was added to the residue, and the mixture was extracted with chloroform. The organic layer was washed with water, dried over anhydrous magnesium sulfate, and the solvent was evaporated. The residue was purified by silica gel column chromatography using n-hexane to obtain first 0.64 g of A isomer of 3-methylthio-1-(1,2,4-triazol-1-yl)-2-[(4-trifluoromethyl)phenyl]butane-2-ol (Compound No. 1) (yield: 9.6%), and then 0.06 g of B isomer (Compound No. 2) (yield: 0.9%).

Example 2

To a solution of 0.35 g (1.06 x 10⁻³ mole) of 3-methylthio-1-(1,2,4-triazol-1-yl)-2-[(4-trifluoromethyl)phenyl]butane-2-ol (Compound No. 1) in 10 ml of chloroform was added under ice-cooling 0.23 g (1.06 x 10⁻³ mole) of m-chloroper-

benzoic acid, and the mixture was stirred at 0 to 10°C for 0.5 hour. The reaction mixture thus obtained was washed with 10% potassium carbonate aqueous solution and then with water, dried over anhydrous magnesium sulfate, and the solvent was evaporated. The residue was purified by silica gel column chromatography using dichloromethane to obtain first 0.23 g of a isomer of 3-methylsulfinyl-1-(1,2,4-triazol-1-yl)-2-[(4-trifluoromethyl)phenyl]butane-2-ol (Compound No. 3) (yield: 62.7%) and subsequently 0.137 g of b isomer (Compound No. 4) (yield: 37.3%).

Example 3

0.30 g (9.1×10^{-4} mole) of 3-methylthio-1-(1,2,4-triazol-1-yl)-2-[(4-trifluoromethyl)phenyl]butane-2-ol (Compound No. 1) was dissolved in 30 ml of chloroform. To the solution was added 0.586 g (2.72×10^{-3} mole) of m-chloroperbenzoic acid, a bit at a time at room temperature, and the mixture was stirred for 2 hours. The reaction mixture thus obtained was washed with 10% potassium carbonate aqueous solution and then with water, dried over anhydrous magnesium sulfate, and the solvent was evaporated. The residue was purified by silica gel column chromatography using dichloromethane to obtain 0.33 g of 3-methylsulfonyl-1-(1,2,4-triazol-1-yl)-2-[(4-trifluoromethyl)phenyl]butane-2-ol (Compound No. 5) (yield: 100%).

Example 4

An oily product of 3-methylsulfonyl-1-(1,2,4-triazol-1-yl)-2-[(4-trifluoromethyl)phenyl]butane-2-ol (Compound No. 6) was prepared by the reaction similar to the reaction of Example 3 from 3-methylthio-1-(1,2,4-triazol-1-yl)-2-[(4-trifluoromethyl)phenyl]butane-2-ol (Compound No. 2).

Example 5

Into 10 ml of t-butanol were dissolved 1.22 g of 2-methylthio-4'-(trifluoromethyl)isobutyrophenone, 1.36 g of trimethylsulfoxonium iodide, 1.07 g of potassium hydroxide, and 0.74 g of 1,2,4-triazole, and the mixture was stirred at 80°C for 6 hours. After the reaction, the solvent was evaporated, water was added to the residue, and the mixture was extracted with chloroform. The organic layer was washed with water, dried over anhydrous magnesium sulfate, and the solvent was evaporated. The residue was purified by silica gel column chromatography using dichloromethane to obtain 0.66 g of an oily product of 3-methyl-3-methylthio-1-(1,2,4-triazol-1-yl)-2-[(4-trifluoromethyl)phenyl]butane-2-ol (Compound No. 7) (yield: 41.1%).

Example 6

To a solution of 0.37 g (1.07×10^{-3} mole) of 3-methyl-3-methylthio-1-(1,2,4-triazol-1-yl)-2-[(4-trifluoromethyl)phenyl]butane-2-ol (Compound No. 7) in 10 ml of chloroform was added under ice-cooling 0.231 g (1.07×10^{-3} mole) of m-chloroperbenzoic acid, and the mixture was stirred at 0 to 10°C for 20 minutes. The reaction mixture thus obtained was washed with 10% potassium carbonate aqueous solution and then with water, dried over anhydrous magnesium sulfate, and the solvent was evaporated. The residue was purified by silica gel column chromatography using dichloromethane to obtain first 0.125 g of a isomer of 3-methyl-3-methylsulfinyl-1-(1,2,4-triazol-1-yl)-2-[(4-trifluoromethyl)phenyl]butane-2-ol (Compound No. 8) (yield: 32.3%), and then 0.155 g of b isomer (Compound No. 9) (yield: 40.0%).

Example 7

0.36 g (1.04×10^{-3} mole) of 3-methyl-3-methylthio-1-(1,2,4-triazol-1-yl)-2-[(4-trifluoromethyl)phenyl]butane-2-ol (Compound No. 7) was dissolved in 40 ml of chloroform. To the solution was added 0.68 g (3.13×10^{-3} mole) of m-chloroperbenzoic acid a bit at a time at room temperature, and the mixture was stirred for 14 hours. The reaction mixture thus obtained was washed with 10% potassium carbonate aqueous solution and then with water, dried over anhydrous magnesium sulfate, and the solvent was evaporated. The residue was purified by silica gel column chromatography using dichloromethane to obtain 0.36 g of 3-methyl-3-methylsulfonyl-1-(1,2,4-triazol-1-yl)-2-[(4-trifluoromethyl)phenyl]butane-2-ol (Compound No. 10) (yield: 91.5%).

Example 8

The following Compounds No. 11-15 were prepared in the same manner as in Examples 1-7.

Compound 11

3-Methylsulfonyl-1-(1,2,4-triazol-1-yl)-2-[(4-trifluoromethyl)phenyl]pentane-2-ol

Compound 12

3-Ethylsulfonyl-3-methyl-1-(1,2,4-triazol-1-yl)-2-[(4-trifluoromethyl)phenyl]butane-2-ol

Compound 13

3-t-Butylsulfonyl-1-(1,2,4-triazol-1-yl)-2-[(4-trifluoromethyl)phenyl]butane-2-ol (a isomer)

Compound 14

3-t-Butylsulfonyl-1-(1,2,4-triazol-1-yl)-2-[(4-trifluoromethyl)phenyl]butane-2-ol (b isomer)

Compound 15

1-[1-(1-methylsulfonyl)cyclopropyl]-2-(1,2,4-triazol-1-yl)-1-[(4-trifluoromethyl)phenyl]ethane-1-ol

Reference Example 8

A mixture of 9.46 g (6.15×10^{-2} mole) of S(-)-methyl-p-tolyl sulfoxide and 100 ml of tetrahydrofuran was added dropwise to 53.3 ml of a cyclohexane solution of lithium diisopropylamide (1.5 mole/l; 7.99×10^{-2} mole) at 5°C and the mixture was stirred at room temperature for 2 hours. To this solution was dropwise added at 10°C 16.1 g (6.15×10^{-2} mole) of 2-methylthio-4'-(trifluoromethyl)isobutyrophenone and the mixture was stirred at room temperature for a further 2 hours. After evaporating the solvent, an oil obtained was treated with 100 ml of saturated NH_4Cl aqueous solution and extracted with ethyl acetate. The organic layer was washed with water, dried over anhydrous magnesium sulfate, and the solvent was evaporated. The residue was submitted to silica gel column chromatography using n-hexane/ethyl acetate to separate two diastereomers (IV-1) and (IV-2) of (-)- α -(1-methyl-1-methylthio)ethyl- α -(4-methylbenzosulfinyl)methyl-4-(trifluoromethyl)benzyl alcohol.

Diastereomer (IV-1)

Amount: 11.19 g (yield: 43.8%), an oily substance

$[\alpha]_D^{25} = -64.2^\circ$ (acetone, $c=0.5$)

IR: ν cm^{-1} 1325, 1065

NMR(CDCl_3) δ ppm: 1.17, 1.31(s, 3H for each), 1.93(s, 3H), 2.43(s, 3H), 3.64(s, 2H), 7.2-8.0(m, 8H)

Diastereomer (IV-2)

Amount: 10.35 g (yield: 40.5%), an oily substance

$[\alpha]_D^{25} = -131.2^\circ$ (acetone, $c=0.5$)

IR: ν cm^{-1} 1330, 1065

NMR(CDCl_3) δ ppm: 1.17, 1.24(s, 3H for each), 2.04(s, 3H), 2.36(s, 3H), 3.50, 4.17(d, 1H for each), 7.0-7.7(m, 8H)

Reference Example 9

11.19 g (2.69×10^{-2} mole) of diastereomer (IV-1) produced in Reference Example 8 was dissolved in 100 ml of dichloromethane. To the mixture was added under ice-cooling 11.6 g (5.38×10^{-2} mole) of 80% m-chloroperbenzoic acid, and the mixture was stirred at 0-10°C for 1 hour. The reaction mixture thus obtained was washed with 20% sodium hydroxide aqueous solution and then with water, dried over anhydrous magnesium sulfate, and the solvent was evaporated. The residue was purified by silica gel column chromatography using chloroform to obtain 10.20 g of crystals of (-)- α -(1-methyl-1-methylsulfonyl)ethyl- α -(4-methylbenzosulfinyl)methyl-4-(trifluoromethyl)benzyl alcohol (V) (yield: 84.6%).

m.p. 100-102°C

$[\alpha]_D^{25} = -35.6^\circ$ (acetone, $c=0.5$)

IR: ν cm^{-1} 1330, 1120, 1065

NMR(CDCl_3) δ ppm: 1.18, 1.39(s, 3H for each), 2.43(s, 3H), 3.19(s, 3H), 3.51, 4.34(d, 1H for each), 7.0-8.2(m, 8H)

Reference Example 10

10.20 g (2.28×10^{-2} mole) of (-)- α -(1-methyl-1-methylsulfonyl)ethyl- α -(4-methylbenzosulfinyl)methyl-4-(trifluoromethyl)benzyl alcohol (V) was dissolved in 50 ml of acetone. To the mixture was added dropwise under ice-cooling 5.36 g (6.83×10^{-2} mole) of acetyl chloride and further added 10.25 g (6.83×10^{-2} mole) of sodium iodide, following which the mixture was stirred at 0-10°C for 1 hour. After the solvent was evaporated, the residue was extracted with ether. The organic layer was washed with water, dried over anhydrous magnesium sulfate, and concentrated. The residue was purified by silica gel column chromatography using n-hexane to obtain 8.93 g of crystals of (+)- α -(1-methyl-1-methylsulfonyl)ethyl- α -[(4-methylphenyl)-thio]methyl-4-(trifluoromethyl)benzyl alcohol (VI) (yield: 90.8%).

m.p. 111-113°C

$[\alpha]_D^{25} = +5.8^\circ$ (acetone, $c=0.5$)

IR: ν cm^{-1} 1330, 1110

NMR(CDCl_3) δ ppm: 1.19, 1.46(s, 3H for each), 2.26(s, 3H), 3.10(s, 3H), 4.02, 4.40(d, 1H for each), 6.9-7.2 (m, 4H), 7.4-7.6(m, 4H)

Reference Example 11

8.93 g (2.07×10^{-2} mole) of (+)- α -(1-methyl-1-methylsulfonyl)ethyl- α -[(4-methylphenyl)-thio]methyl-4-(trifluoromethyl)benzyl alcohol (VI) was dissolved in 50 ml of dichloromethane, and to the mixture was added dropwise under ice-cooling 100 ml of a dichloromethane solution of triethyloxonium tetrafluoroborate (1 mole/l; 0.1 mole), followed by stirring at room temperature overnight. After an addition of 40 ml of 20% sodium hydroxide aqueous solution under ice-cooling, the mixture was stirred at room temperature for a further 1 hour. The organic layer was separated, washed with water, dried over anhydrous magnesium sulfate, and the solvent was evaporated. The residue was purified by silica gel column chromatography using n-hexane to obtain 5.45 g of an oily product of (-)-2-(1-methyl-1-methylsulfonyl)ethyl-2-[(4-(trifluoromethyl)phenyl)oxy]lane (VII) (yield: 85.6%).

$[\alpha]_D^{25} = -20.6^\circ$ (acetone, $c=0.5$)

IR: cm^{-1} 1325, 1110

NMR(CDCl_3) δ ppm: 1.47, 1.52(s, 3H for each), 2.81, 3.62 (d, 1H for each), 2.89(s, 3H), 7.61(s, 4H)


Example 9

1.50 g (4.87×10^{-3} mole) of (-)-2-(1-methyl-1-methylsulfonyl)ethyl-2-[(4-(trifluoromethyl)phenyl)oxy]lane (VII) and 1.73 g (2.5×10^{-2} mole) of triazole were dissolved in 45 ml of dimethylformamide. To this solution was added 0.69 g (5.0×10^{-3} mole) of potassium carbonate and the mixture was stirred at room temperature overnight. The resultant mixture was dissolved in a water/ether mixture. The organic layer was separated, washed with water, dried over anhydrous magnesium sulfate, and the solvent was evaporated. The residue was purified by silica gel column chromatography using chloroform. 1.198 g of crystals of (-)-3-methyl-3-methylsulfonyl-1-(1,2,4-triazol-1-yl)-2-[(4-trifluoromethyl)phenyl]butane-2-ol (Compound No. 16) was obtained by recrystallization from an isopropylether/ diethylether mixed solvent (yield: 54.2%).

$[\alpha]_D^{25} = -9.6^\circ$ (acetone, $c=0.25$)

Characteristics, IR, and NMR of compounds prepared in Examples 1-9 are shown in Table 2.

TABLE 2



Compound No.	R ₁	R ₂	R ₃	n	Characteristics m.p. (°C)	IR (cm ⁻¹)	NMR (CDCl ₃) δ (ppm)
1	CH ₃	H	CH ₃	0	White crystals 97-100	3150, 1330	1.21(d, 3H), 2.16(s, 3H), 3.02(q, 1H), 4.63, 5.00(d, 1H for each), 7.42, 7.54 (d, 2H for each), 7.81(s, 2H)
2	CH ₃	H	CH ₃	0	Oil	3220, 1325	1.28(d, 3H), 1.93(s, 3H), 2.84(q, 1H), 4.80, 4.87(d, 1H for each), 7.60(s, 4H), 7.88, 8.28 (s, 1H for each)
3	CH ₃	H	CH ₃	1	White crystals 147-150	3180, 1325	1.20(d, 3H), 2.64(s, 3H), 2.96(q, 1H), 4.52, 5.09(d, 1H for each), 7.64(s, 4H), 7.94, 8.26(s, 1H for each)
4	CH ₃	H	CH ₃	1	White crystals 168-169	3200, 1325	1.21 (d, 3H), 2.66(s, 3H), 3.28(q, 1H), 4.80, 5.04(d, 1H for each), 7.36, 7.60 (d, 2H for each), 7.94, 8.36(s, 1H for each)

TABLE 2 (Continued)

Compound No.	R ₁	R ₂	R ₃	n	Characteristics m.p. (°C)	IR (ν _{cm⁻¹)}	NMR (CDCl ₃) δ (ppm)
5	CH ₃	H	CH ₃	2	White crystals 172-175	3350, 1320	1.30 (d, 3H), 3.06(s, 3H), 3.38(q, 1H), 5.26, 5.39(d, 1H for each), 7.44, 7.58 (d, 2H for each), 7.71, 7.80(s, 1H for each)
6	CH ₃	H	CH ₃	2	Oil	3400, 1320	1.54 (d, 3H), 2.80(s, 3H), 3.2-3.6(m, 1H), 4.90, 5.10(d, 1H for each), 7.59(s, 4H), 7.77, 8.13 (s, 1H for each)
7	CH ₃	CH ₃	CH ₃	0	Oil	3400, 1320	1.31, 1.38(s, 3H for each), 2.02(s, 3H), 4.88, 5.14(d, 1H for each), 7.50, 7.64 (d, 2H for each), 7.71, 7.99 (s, 1H for each)
8	CH ₃	CH ₃	CH ₃	1	White crystals 159-162	3300, 1325	1.05, 1.55(s, 3H for each), 2.49(s, 3H), 4.76, 5.56(d, 1H for each), 7.54, 7.68 (d, 2H for each), 7.73, 8.15(s, 1H for each)
9	CH ₃	CH ₃	CH ₃	1	White crystals 82-85	3250, 1330	1.07, 1.47(s, 3H for each), 2.59(s, 3H), 5.04(s, 2H), 7.58 (s, 4H), 7.74, 8.21(s, 1H for each)
10	CH ₃	CH ₃	CH ₃	2	White crystals 156-157	3150, 1325	1.27, 1.53(s, 3H for each), 3.21(s, 3H), 4.96, 5.20(d, 1H for each), 7.4-7.7 (m, 4H), 7.69, 7.96(s, 1H for each)

TABLE 2 (Continued)


Compound No.	R ₁	R ₂	R ₃	n	Characteristics m.p. (°C)	IR (ν _{cm⁻¹})	NMR (CDCl ₃) δ (ppm)
11	C ₂ H ₅	H	CH ₃	2	Oil	3400, 1320	0.74(t, 3H), 0.8-2.0(m, 3H), 3.10(s, 3H), 4.83, 5.30(d, 1H for each), 7.4-8.0 (m, 6H).
12	CH ₃	CH ₃	C ₂ H ₅	2	White crystals 169-170	3150, 1325	1.27, 1.54(s, 3H for each), 1.49(t, 3H), 3.1-3.8(m, 2H), 5.17, 5.43(d, 1H for each), 7.4-7.8(m, 4H), 7.66, 7.99(s, 1H for each)
13	CH ₃	H	t-C ₄ H ₉	2	White crystals 152-153	3350, 1320	1.35(s, 9H), 1.59(d, 3H), 3.60(q, 1H), 4.55, 4.99(d, 1H for each), 7.62(s, 4H), 7.91, 8.27(s, 1H for each)
14	CH ₃	H	t-C ₄ H ₉	2	White crystals 211-212	3400, 1320	1.35(s, 9H), 1.36(d, 3H), 3.50(q, 1H), 4.78, 5.23(d, 1H for each), 7.61, 7.74 (d, 2H for each), 7.87, 8.21(s, 1H for each)
15	-CH ₂ CH ₂ -	CH ₃	CH ₃	2	White crystals 164-165	3200, 1320	0.7-1.6(m, 4H), 2.25(s, 3H), 5.34(s, 2H), 7.65, 7.74(d, 2H for each), 7.91, 8.53 (s, 1H for each)
16	CH ₃	CH ₃	CH ₃	2	White crystals 97-98	1330, 1110	1.26, 1.53(s, 3H for each), 3.21(s, 3H), 5.13, 5.47(d, 1H for each), 7.4-7.8 (m, 4H), 7.67, 7.99(s, 1H for each)

Obviously, numerous modifications and variations of the present invention are possible in light of the above teachings. It is therefore to be understood that within the scope of the appended claims, the invention may be practiced otherwise than as specifically described herein.

Claims

Claims for the following Contracting States : BE, CH, DE, FR, GB, IT, LI, NL, SE

1. A triazole derivative of formula (I),




wherein R^1 and R^2 represent a hydrogen atom or a C_{1-6} -alkyl group, or R^1 and R^2 together form an ethylene, trimethylene or tetramethylene group, R^3 represents a C_{1-6} -alkyl group, and n denotes an integer of 0 to 2, provided that R^1 and R^2 are not both a hydrogen atom at the same time, and a salt thereof.

2. A composition for curing mycoses comprising an effective amount of a triazole derivative or a salt thereof according to Claim 1 and a pharmaceutically acceptable carrier.


Claim for the following Contracting State : ES

1. A process for the preparation of a triazole derivative of formula (I),




wherein R^1 and R^2 represent a hydrogen atom or a C_{1-6} -alkyl group, or R^1 and R^2 together form an ethylene, trimethylene or tetramethylene group, R^3 represents a C_{1-6} -alkyl group, and n denotes an integer of 0 to 2, provided that R^1 and R^2 are not both a hydrogen atom at the same time, and a salt thereof, which process comprises the steps of:

a) epoxylating a compound of formula (II)





wherein R^1 , R^2 , and R^3 are as defined above, and reacting the obtained product with triazole to obtain a product of formula I-1



wherein R^1 , R^2 and R^3 are as defined above;

b) optionally oxidizing the compound of formula I-1 to obtain a compound of formula 1-2 or 1-3






15 wherein R^1 , R^2 and R^3 are as defined above.

Patentansprüche

20 Patentansprüche für folgende Vertragsstaaten : BE, CH, DE, FR, GB, IT, LI, NL, SE

1. Triazolderivat der Formel (I)




40 wobei R^1 und R^2 für ein Wasserstoffatom oder eine C_{1-6} -Alkylgruppe stehen, oder R^1 und R^2 zusammen eine Ethylen-, Trimethylen- oder Tetramethylengruppe bilden, R^3 für eine C_{1-6} -Alkylgruppe steht, und n für eine ganze Zahl von 0 bis 2 steht, mit der Maßgabe, daß R^1 und R^2 nicht beide gleichzeitig für ein Wasserstoffatom stehen, sowie ein Salz davon.

45 2. Mittel zur Heilung von Mykosen, das eine wirksame Menge eines Triazolderivats oder eines Salzes davon nach Anspruch 1, sowie einen pharmazeutisch unbedenklichen Träger umfaßt.


Patentanspruch für folgenden Vertragsstaat : ES

50 1. Verfahren zur Herstellung eines Triazolderivats der Formel (I)




15 wobei R¹ und R² für ein Wasserstoffatom oder eine C₁₋₆-Alkylgruppe stehen, oder R¹ und R² zusammen eine Ethylen-, Trimethylen- oder Tetramethylen-Gruppe bilden, R³ für eine C₁₋₆-Alkylgruppe steht, und n für eine ganze Zahl von 0 bis 2 steht, mit der Maßgabe, daß R¹ und R² nicht beide gleichzeitig für ein Wasserstoffatom stehen, sowie ein Salz davon, wobei das Verfahren folgende Schritte umfaßt:

20 (a) Epoxidierung einer Verbindung der Formel (II)





35 wobei R¹, R² und R³ wie oben definiert sind, und Umsetzung des erhaltenen Produkts mit Triazol, um ein Produkt der Formel I-1 zu schaffen



50 wobei R¹, R² und R³ wie oben definiert sind;

(b) gegebenenfalls Oxidation der Verbindung der Formel I-1, um eine Verbindung der Formel I-2 oder I-3 zu schaffen




wobei R^1 , R^2 und R^3 wie oben definiert sind.

Revendications

Revendications pour les Etats contractants suivants : BE, CH, DE, FR, GB, IT, LI, NL, SE

1. Dérivé du triazole de formule (I)




dans laquelle R^1 et R^2 représentent un atome d'hydrogène ou un groupe alkyle en C_{1-6} , ou R^1 et R^2 forment ensemble un groupe éthylène, triméthylène ou tétraméthylène, R^3 représente un groupe alkyle en C_{1-6} et n désigne un entier de 0 à 2, à la condition que R^1 et R^2 ne soient pas tous les deux un atome d'hydrogène en même temps, et un sel de celui-ci.

2. Composition pour guérir les mycoses, comprenant une quantité efficace d'un dérivé du triazole ou d'un sel de celui-ci, selon la revendication 1, et un excipient acceptable d'un point de vue pharmaceutique.


R v ndication pour l'Etat contractant suivant : ES

1. Procédé pour préparer un dérivé du triazole de formule (I)




dans laquelle R^1 et R^2 représentent un atome d'hydrogène ou un groupe alkyle en C_{1-6} , ou R^1 et R^2 forment ensemble un groupe éthylène, triméthylène ou tétraméthylène, R^3 représente un groupe alkyle en C_{1-6} et n désigne un entier de 0 à 2, à la condition que R^1 et R^2 ne soient pas tous les deux un atome d'hydrogène en même temps, et un sel de celui-ci, lequel procédé comprend les étapes :

a) d'époxydation d'un composé de formule (II)





dans laquelle R^1 , R^2 et R^3 sont tels que définis ci-dessus, et de réaction du produit obtenu, avec du triazole pour donner un produit de formule I-1



où R^1 , R^2 et R^3 sont tels que définis ci-dessus ;

b) éventuellement, d'oxydation du composé de formule I-1 pour donner un composé de formule I-2 ou I-3



25

où R¹, R² et R³ sont tels que définis ci-dessus.

THIS PAGE BLANK (USPTO)